## **OCD Repair or OATs post op Rehab**

## **Post Operative Weeks 1-6:**

- Home Exercise Program
  - o Remain non weight bearing
  - o Range of motion as tolerated

## Weeks 6-14: Begin formal Physical Therapy

- Goals
  - o Restore full ROM
  - o Restore normal Gait
  - o Demonstrate ability to ascend and descend stairs with good control, and without pain
  - o Independent with home exercise program
- Precautions
  - o Avoid descending stairs reciprocally until adequate quad control and proper alignment
  - o Avoid pain with ther ex, and functional activities
  - Avoid running and other impact or sporting activities
- Exercises
  - o Progress weight bearing as quad control allows, D/C crutches when gait is non-antalgic
  - Active assisted range of motion exercises
  - SLR in all planes
  - o Balance and other neuromuscular training
  - Stationary bike
  - Leg press, lower extremity PRE's
  - Squat 0-60°
  - Stair master / elliptical machines
  - Forward step up/down

## Goals:

- o Demonstrate ability to run pain free
- o Maximize strength and flexibility to meet demands of ADL's
- o Independence with gym program for maintenance and progression
- Exercises: Continue LE strengthening as previous
  - o Progress squats to 90°
  - o Lunges
  - o Begin forward jogging on treadmill (4 months post op)
  - o Begin agility / sport specific specific program (when strength base is sufficient)
  - o Begin plyometric program (when strength base is sufficient)